

CONTENTS

Letter to the Editor

- Robot-assisted pulmonary lobectomy**
Sami Karapolat, Suat Gezer 60

Clinical Studies

- The most encountered groups of genetic disorders in Giza Governorate, Egypt**
Afifi HH, El-Ruby MO, El-Bassyouni HT, Ismail SI, Aglan MS, El-Harouni AA, Mazen IM, Zaki MS, Bassiouni RI, Hosny LA, El Kamah GY, El Kotoury AI, Ashour AM, Abdel-Salam GM, El-Gammal MA, Hamed K, Kamal RM, El-Nekhely I, Temtamy SA 62

This study presents the prevalence, relative frequency, and analysis of genetic diseases/malformations in 73 260 individuals, from Giza governorate, Egypt. The overall parental consanguinity rate among the 3417 diagnosed cases was 55 %, ranging from 29.5–75 %. 3 % of neonates in NICU had genetic or congenital disorders.

- Plasma homocysteine level and 677C→T mutation on the MTHFR gene in patients with venous thromboembolism**
Hanta I, Soydas Y, Karatasli M, Koseoglu Z, Satar S, Hasturk S 70

The aim of this study was to evaluate the role of plasma total homocysteine level and 677C→T mutation on the methylenetetrahydrofolate reductase (MTHFR) gene in the development of venous thromboembolism. It is thought that a high plasma total homocysteine may cause venous thromboembolism.

- Real-life experiences in migraine therapy**
Turcani P, Farkasova D, Marcek T, Turcani M 74
- The results of this open-label study reflective of real-life clinical practice provide further evidence for the efficacy, consistency of response, safety and tolerability of oral eletriptan in acute treatment of migraine headache, associated symptoms as well as functional impairment.

- Metabolic abnormalities in patients with recurrent stone formation in a hot territory**
Nader Nouri-Majalan, Behnam Baghianimoghadam, Noushin Amiri, Sara Moghaddasi-Moosavi 79

The aim of this study is to evaluate the metabolic abnormalities in patients with recurrent stone formation in a hot territory in our country. The findings suggest that lower urine volume has an important role in nephrolithiasis in hot areas. Higher urine sodium and uric acid and lower serum phosphor correlate with higher urine calcium in stone formers.

- Retrograde endarterectomy of iliac arteries**
Beno M, Rumenapf G 83
- Authors retrospectively analyze a group of patients with symptomatic peripheral arterial occlusive disease (PAOD) who underwent an arterial reconstruction with a remote retrograde endarterectomy combined with intraoperative angiography and endovascular techniques as a hybrid procedure. The aim of this study was to analyze complications of this procedure.

- Cryothermal ablation of typical atrial flutter**
Misikova S, Stancak B 90
- An alternative source of energy for the ablation of cardiac tachyarrhythmias is cryoenergy. Efficacy and safety profile of cryoablation are similar with RF ablation. Procedural time in the cryoablation group is significantly longer but the use of analgesics can be completely avoided.

- Phacoemulsification of mature and hard nuclear cataracts**
Ilavska M, Kardos L 93

The aim of the study was to compare the best corrected visual acuity (BCVA) before and on the 1st day after the cataract surgery for mature or hard nuclear cataract. The phacoprobe Neo-SoniX equipment Legacy (Alcon Laboratories) was used for the cataract surgery. Trypan blue was used for vital staining of the anterior capsule in case of mature or white cataract. The prevalence and the effect of the complication on CVA were evaluated.

- Digital and manual cephalometric analyses**
Thurzo A, Javorka V, Stanko P, Lysy J, Suchancová B, Lehotska V, Valkovic L, Makovnik M 97
- Evaluation of 100 analog cephalograms digitized by digital camera and analyzed by Dolphin Imaging 10 compared the manual cephalometric tracing and the computer method. The results show that software analysis can substitute the manual method with even better accuracy, as well as that digital cameras can be used for analog x-ray digitalization.

Case Reports

- Gait apraxia**
Mihalj M, Titlic M, Marovic A, Bulovic B, Srdelic-Mihalj S 101

Gait apraxia is most commonly a part of the Hakimov triad (gait apraxia, urinary incontinence, dementia) in normotensive hydrocephalus (NPH), although it may be a symptom of some other conditions. It remains a diagnostic and therapeutic challenge, especially in cases where NPH is complicated with parkinsonism. Many ancillary investigations have been described that can increase the probability of selecting the appropriate candidates for a shunt.

- Rectal prolapse**
Sekac J, Labas P, Skultety J, Prochotsky A, Skubla R, Okolicany R 103

It is not so much the diagnosis that offers the surgeon a wide range of opportunities in the technical solutions of rectal prolapse. Currently there are at least 130 different techniques used in the surgical treatment of rectal prolapse and in fact none of these procedures has been shown most effective for any one patient. All of these surgical techniques have one denominator in common, namely that although the risk of recurrence varies with authors it seldom exceeds 20 %.

- A very rare localisation of pilomatixoma**
Samet Yalcin, Burak Veli Ulger, Omer Parlak, Cagatay Sisman, Aykut Onurseven, Ahmet Kusdemir 108

Pilomatixoma is a rare skin neoplasm which is usually localized on face, neck and lower extremities and its etiology is still unknown. We report a case of a 32-year-old male patient with a rare localisation of pilomatixoma on the right shoulder. The presumed diagnosis was sebaceous cyst. Excision biopsy was performed.